

- b) Find the directional derivative of $\phi = xy + yz + zx$ in the direction of the vector $i + 2j + 2k$ at the point $(1,2,0)$.
- c) If $r = xi + yj + zk$ then show that $\text{grad } r^n = nr^{n-2}r$.
- d) Verify Stoke's theorem for $F = (x^2 + y^2)i - 2xyj$ taken round the rectangle bounded by $x = \pm a, y = 0, y = b$.

OR

Prove that $\text{div. grad } r^m = \nabla \cdot \nabla r^m = m(m+1)r^{m-2}$

Roll No

BE - 301**B.E. III Semester**

Examination, June 2015

Engineering Mathematics - II

(Common for all Branches)

Time : Three Hours**Maximum Marks : 70**

- Note:** i) Answer five questions. In each question part A, B, C is compulsory and D part has internal choice.
- ii) All parts of each questions are to be attempted at one place.
- iii) All questions carry equal marks, out of which part A and B (Max.50 words) carry 2 marks, part C (Max.100 words) carry 3 marks, part D (Max.400 words) carry 7 marks.
- iv) Except numericals, Derivation, Design and Drawing etc.

UNIT - I

1. a) What is periodic function, give an example of periodic function?
- b) What are the dirichlet's condition for a Fourier expansion?
- c) Find the Fourier transform of e^{-ax^2} , where $a > 0$.
- d) Expand $f(x) = x \sin x$, $0 < x < 2\pi$ in a Fourier series.

[2]

OR

Express $f(x) = x$ as a

- i) Half range cosine series in $0 < x < 2$
 ii) Half range sine series in $0 < x < 2$

UNIT - II

2. a) Find $L\{2\sin t \cos t\}$.
 b) Write the Linearly property of Laplace Transform.
 c) Find the Laplace Transform of $f(t)$,

$$\text{where } f(t) = \begin{cases} 2, & 0 \leq t \leq 2 \\ t-1, & 2 \leq t \leq 3 \\ 7, & t > 3 \end{cases}$$

d) Evaluate $L^{-1}\left\{\frac{6s^2 + 22s + 16}{s^3 + 6s^2 + 11s + 6}\right\}$.

OR

Using Convolution Theorem, Evaluate $L^{-1}\left\{\frac{s}{(s^2 + a^2)^2}\right\}$.

UNIT - III

3. a) Explain the ordinary point and singular point of differential equation.
 b) Give the complete solution of differential equation when the roots of indicial equations are equal.

[3]

- c) Solve the differential equation

$$\frac{d^2y}{dx^2} - 2 \tan x \frac{dy}{dx} + 5y = \sec x.e^x$$

- d) Solve $(1-x^2)\frac{d^2y}{dx^2} + 2x\frac{dy}{dx} + y = 0$ in series solution.

OR

Solve by method of variation of parameter $(D^2+1)y = x$.

UNIT - IV

4. a) Find the Partial differential equation by eliminating a and b from the relation $(x-a)^2 + (y-b)^2 = z^2 - c$.
 b) Solve $yzp + zxq = xy$.
 c) Solve $\frac{\partial^2 z}{\partial x^2} + 2\frac{\partial^2 z}{\partial x \partial y} + \frac{\partial^2 z}{\partial y^2} = e^{3x+2y}$.
 d) Solve by Charpits method $(p^2 + q^2)y = qz$.

OR

Solve $\frac{\partial u}{\partial x} = 2\frac{\partial u}{\partial t} + u$ by the method of separation of variables. Where $u(x,0) = 6e^{-3x}$.

UNIT - V

5. a) If $u = t^2i - tj + (2t+1)k$
 $v = (2t-3)i + j - tk$
 find $\frac{d}{dt}(u.v)$? at $t = 1$.